

THE FUNCTIONAL INTEGRAL FOR A FREE
PARTICLE ON A HALF-PLANE*

Michel Carreau

Boston University
Department of Physics
590 Commonwealth Avenue
Boston, Massachusetts 02215 U.S.A.
E-mail:CARREAU@BUPHY.BU.EDU

To be published in : *Journal of Mathematical Physics*

BU-HEP-91-23

Typeset in $T_{\text{E}}X$

November 1991

* This work is supported in part by funds provided by the U. S. Department of Energy (D.O.E.) under contract #DE-AC02-76ER03069, and by the Fonds pour la Formation de Chercheurs et l'Aide à la Recherche.

ABSTRACT

A free non-relativistic particle moving in two dimensions on a half-plane can be described by self-adjoint Hamiltonians characterized by boundary conditions imposed on the systems. The most general boundary condition is parameterized in terms of the elements of an infinite-dimensional matrix. We construct the Brownian functional integral for each of these self-adjoint Hamiltonians. Non-local boundary conditions are implemented by allowing the paths striking the boundary to jump to other locations on the boundary. Analytic continuation in time results in the Green's functions of the Schrödinger equation satisfying the boundary condition characterizing the self-adjoint Hamiltonian.

I. INTRODUCTION

In a paper by Clark et al., and later on by Gaveau et al. and Farhi et al. (see Ref. [1]) the functional integral for a particle moving on a half-line was derived. It was found that the measure for the functional integral depends upon the one parameter family of boundary conditions which guarantee that probability does not leave the half-line. Farhi, Gutmann and the present author² derived the functional integral for a free particle moving in a box in 1-D for the system subject to a four-parameter family of boundary conditions consistent with unitarity. They found that the measure for the functional integral depends upon these four parameters and that the paths are allowed to jump from wall to wall.

In this paper, we show how to generalize these ideas to more than one dimension. In Section II, we describe the most general boundary condition leading to unitary time evolution for the free particle moving on a half-plane. (Equivalently, we find the self-adjoint extensions of the free Hamiltonian.) In Section III, we construct the Brownian functional integral representation of the Brownian Green's function corresponding to each self-adjoint Hamiltonian. The Green's function of the Schrödinger equation are obtained by an analytic continuation of the Brownian Green's function. We point out how to generalize these results to more complex quantum systems. In the Appendix we give the functional integrals of a one dimensional quantum systems made out of any number of half-lines joined together at their origin.

II. PARTICLE ON A HALF-PLANE — SELF-ADJOINT EXTENSIONS

Consider the motion of a free particle on a half-plane in two dimensions ($-\infty < y < +\infty$ and $x > 0$). The Hilbert space consists of all square integrable functions on the half-plane. The domain of the Hamiltonian

$$\hat{H} = -\frac{1}{2} \frac{d^2}{dx^2} - \frac{1}{2} \frac{d^2}{dy^2} \quad (2.1)$$

must be chosen so that $\hat{H}$ is Hermitian, which is the condition

$$\int_{-\infty}^{+\infty} \int_0^{\infty} \psi^* (\hat{H} \phi) dx dy = \int_{-\infty}^{+\infty} \int_0^{\infty} (\hat{H} \psi)^* \phi dx dy \quad (2.2)$$

or equivalently

$$\int_{-\infty}^{+\infty} dy \left[\psi^* \frac{d\phi}{dx} - \frac{d\psi^*}{dx} \phi \right] \Big|_{x=0} = 0 \quad (2.3)$$

for all ψ and ϕ in the domain of $\hat{H}$.

The condition (2.3) can be realized most generally if for every χ in the domain of $\hat{H}$

$$\frac{d\chi}{dx}(x, y) \Big|_{x=0} = \int_{-\infty}^{+\infty} dy' g(y', y) \chi(0, y') \quad (2.4)$$

for each y , with g an arbitrary function obeying

$$g(y', y) = g^*(y, y') \quad (2.5)$$

A similar result was obtained in Ref. [3]. Let us pick the following parameterization for $g(y', y)$

$$g(y', y) = \left\{ [\rho(y) + \alpha(y)] \delta(y' - y) - \rho(y') f(y', y) e^{ih(y', y)} \right\} \quad (2.6)$$

where δ is the delta-function and α, h, ρ, f are arbitrary smooth real functions with the following restrictions: ρ, f are non-negative, h is antisymmetric in its arguments,

$$\rho(y) f(y, y') = \rho(y') f(y', y) \quad (2.7)$$

and

$$\int_{-\infty}^{+\infty} dy f(y', y) = 1 \quad . \quad (2.8)$$

We can show that $g(y', y)$ uniquely determines the functions α, h, ρ, f . Given a g satisfying (2.5) if we choose the domain of $\hat{H}$ to be functions obeying (2.4) then $\hat{H}$ is self-adjoint and probability does not leave the half-plane. We see that we have infinitely many self-adjoint extensions indexed by the function $g(y', y)$. To each self-adjoint extension there corresponds a unique Green's function $G_g(\vec{z}_i, \vec{z}_f, t)$ with $\vec{z}_a = (x_a, y_a)$, $a = i, f$ on the half-plane. The Green's function can be expressed as

$$G_g(\vec{z}_i, \vec{z}_f, t) = \langle \vec{z}_f | e^{-i\hat{H}t} | \vec{z}_i \rangle_g \quad . \quad (2.9)$$

Thought of as a function of $\vec{z}_f$, G_g satisfies the Schrödinger equation and the boundary condition (2.4), that is

$$\left. \frac{d}{dx_f} G_g(\vec{z}_i, \vec{z}_f, t) \right|_{x_f=0} = \int dy'_f g(y'_f, y_f) G_g(\vec{z}_i, (0, y'_f), t) \quad . \quad (2.10)$$

If the boundary condition (2.4) has the property that when it is satisfied by $\chi(x, y)$ it is also satisfied by $\chi(x, y + y_0)$ for an arbitrary y_0 , we say it is translationally invariant in the y -direction. If we want a translationally invariant boundary condition in the y -direction, then $g(y, y')$ can depend only on $y - y'$.

Condition (2.4) says that the *net* current flowing out of the half-plane is zero. The condition that the current flow at each point on the edge of the half-plane be zero is that the integrand in (2.3) vanishes for each y . This can be realized by taking $\rho(y) = 0$, leading to

$$g(y', y) = \alpha(y) \delta(y' - y) \quad , \quad (2.11)$$

the most general *local* boundary condition. If we want to enforce the translation invariance in the y -direction and the condition that no current flow at each y , we take $\alpha(y)$ independent of y . Then g is simply

$$g(y', y) = \alpha \delta(y' - y) \quad . \quad (2.12)$$

With these restrictions there is only a one-parameter family of self-adjoint extensions.

III. PARTICLE ON A HALF-PLANE — FUNCTIONAL INTEGRAL

In this section, we construct the Brownian functional integrals corresponding to the self-adjoint extensions of the free Hamiltonian constructed in the previous section. The analytic continuation of the Brownian functional integrals results in the Green's functions of the Schrödinger equation.

1. Preliminaries

In order to have a mathematically well-defined measure on a class of paths, we work with Brownian functional integrals. On the whole plane the Brownian functional integral

$$\mathcal{A}_{\text{wp}}^{\text{B}}(\vec{z}_i, \vec{z}_f, \tau) = \int [d\vec{q}_{\text{wp}}] e^{-\frac{1}{2} \int_0^\tau \dot{\vec{q}}_{\text{wp}}^2 d\tau'} = \frac{1}{2\pi\tau} e^{-|\vec{z}_f - \vec{z}_i|^2 / 2\tau} \quad (3.1)$$

with $\vec{q}_{\text{wp}}(0) = \vec{z}_i$ and $\vec{q}_{\text{wp}}(\tau) = \vec{z}_f$, is the Green's function of the Schrödinger equation with imaginary time $t = -i\tau$. We call $\mathcal{A}_{\text{wp}}^{\text{B}}$ the Brownian Green's function. The symbol $[d\vec{q}_{\text{wp}}] e^{-\frac{1}{2} \int_0^\tau \dot{\vec{q}}_{\text{wp}}^2 d\tau'}$ is to be interpreted as the Brownian measure on paths $\vec{q}_{\text{wp}}$. This measure gives zero weight to discontinuous paths. The Green's function, G_{wp} , of the Schrödinger equation on the whole plane is obtained by the analytic continuation of $\mathcal{A}_{\text{wp}}^{\text{B}}$:

$$G_{\text{wp}}(\vec{z}_i, \vec{z}_f, t) \equiv \mathcal{A}_{\text{wp}}^{\text{B}}(\vec{z}_i, \vec{z}_f, it) \quad (3.2)$$

When working on the half-plane, we use the *reflecting measure*, which is induced by taking the absolute value of the x -component of whole plane Brownian motion. That is, if $\vec{q}_{\text{wp}} = (q_{1\text{wp}}, q_{2\text{wp}})$ is a whole plane path then $\vec{q}^* \equiv (|q_{1\text{wp}}|, q_{2\text{wp}}) \equiv (q_x^*, q_y)$ lies on the half-plane and the probability distribution, or measure, on $\vec{q}_{\text{wp}}$ directly induces a probability distribution on $\vec{q}^*$. For each path $\vec{q}^* = (q_x^*, q_y)$ which goes from $\vec{z}_i$ to $\vec{z}_f$ on the half-plane in time τ , there exists a “local time” which tells us how much time the path spends near the boundary of the half-plane ($x \approx 0$).

$$\ell(\tau) \equiv \lim_{\epsilon \rightarrow 0} \frac{1}{2\epsilon} \int_0^\tau I(q_x^*(\tau') < \epsilon) d\tau' \quad (3.3)$$

where $I(A) = 1$ or 0 according to whether A occurs or not. (More informally, we can think of $\ell(\tau)$ as $\int_0^\tau \delta(q_x^*(\tau')) d\tau'$.)

We now define the half-plane Brownian functional integral,

$$\mathcal{A}_r^{\text{B}}(\vec{z}_i, \vec{z}_f, \tau) = \int [d\vec{q}^*] e^{-\frac{1}{2} \int_0^\tau \dot{\vec{q}}^{*2} d\tau'} e^{-r\ell(\tau)} \quad (3.4)$$

where r is a real parameter and the functional integral is over all half-plane reflected paths $\vec{q}^* = (q_x^*, q_y)$ from $\vec{z}_i$ to $\vec{z}_f$ in time τ . The $[d\vec{q}^*]$ times the first exponential is to be interpreted as the reflected Brownian measure. There is a Brownian motion theorem⁴ that the limit in (3.3) exists for almost every path so (3.4) is sensibly defined. The Brownian motion on the half-plane, $\vec{q}^*(\tau')$, is the Cartesian product of a reflecting Brownian motion in the x -direction, $q_x^*(\tau')$, and a whole line Brownian motion in the y -direction, $q_y(\tau')$. Since these

motions are completely independent, we can write the reflecting measure on the half-plane as the product of the reflecting measure in the x -direction times the whole line measure in the y -direction. Moreover, the local time, $\ell(\tau)$, depends only on the motion in the x -direction. Therefore, the functional integral (3.4) factors as follows

$$\begin{aligned} \mathcal{A}_r^B(\vec{z}_i, \vec{z}_f, \tau) &= \mathcal{A}_{x,r}^B(x_i, x_f, \tau) \mathcal{A}_y^B(y_i, y_f, \tau) \\ &\equiv \left\{ \int [dq_x^*] e^{-\frac{1}{2} \int_0^\tau \dot{q}_x^{*2} d\tau'} e^{-r\ell(\tau)} \right\} \left\{ \int [dq_y] e^{-\frac{1}{2} \int_0^\tau \dot{q}_y^2 d\tau'} \right\} \end{aligned} \quad (3.5)$$

where $\vec{z}_a = (x_a, y_a)$, for $a = i, f$. It follows that $\mathcal{A}_r^B$ satisfies the boundary condition

$$\left. \frac{d}{dx_f} \mathcal{A}_r^B(\vec{z}_i, \vec{z}_f, \tau) \right|_{x_f=0} = r \mathcal{A}_r^B(\vec{z}_i, \vec{z}_f, \tau) \Big|_{x_f=0} \quad (3.6)$$

since $A_{x,r}^B$ has been shown¹ to satisfy the same boundary condition. The analytic continuation, $G_r(\vec{z}_i, \vec{z}_f, t) \equiv \mathcal{A}_r^B(\vec{z}_i, \vec{z}_f, it)$ is the Green's function of the Schrödinger equation for a free particle on the half-plane satisfying (3.6). This one-parameter family of Green's functions corresponds to the one-parameter family of self-adjoint Hamiltonians defined in the last section by (2.12), i.e. those with translationally invariant local boundary conditions.

2. General Local Boundary Condition

In this subsection, we build the Brownian functional integrals for the self-adjoint extensions corresponding to the boundary condition (2.4) with g given by (2.11). Let us generalize the term, $r\ell(\tau)$, in the functional integral (3.4) to

$$\int_0^\tau \alpha(q_y(\tau')) \dot{\ell}(\tau') d\tau' \quad (3.7)$$

for a path $\vec{q}^* = (q_x^*, q_y)$ with local time ℓ , where $\dot{\ell} \equiv d\ell/d\tau'$ and α is an arbitrary real function. The factor $\dot{\ell}$ gives a contribution different from zero whenever the paths hit the edge of the half-plane. More informally, we can think of (3.7) as

$$\int_0^\tau \int_{-\infty}^{+\infty} \alpha(y_0) \delta(q_y(\tau') - y_0) \delta(q_x^*(\tau')) dy_0 d\tau' , \quad (3.8)$$

a continuous sum of delta-function potentials along the edge of the half-plane with different strengths for different y . The functional integral on the half-plane becomes

$$\mathcal{A}_\alpha^B(\vec{z}_i, \vec{z}_f, \tau) = \int [d\vec{q}^*] e^{-\frac{1}{2} \int_0^\tau \dot{\vec{q}}^{*2} d\tau' - \int_0^\tau \alpha(q_y(\tau')) \dot{\ell}(\tau') d\tau'} . \quad (3.9)$$

The expression (3.7) exists for almost every path^{5,6} so (3.9) is sensibly defined. In the next subsection we will show that the analytic continuation $G_\alpha(\vec{z}_i, \vec{z}_f, t) \equiv \mathcal{A}_\alpha^B(\vec{z}_i, \vec{z}_f, it)$ is the Green's function of the Schrödinger equation satisfying the most general local boundary condition

$$\left. \frac{d}{dx_f} G_\alpha(\vec{z}_i, \vec{z}_f, t) \right|_{x_f=0} = \alpha(y_f) G_\alpha(\vec{z}_i, \vec{z}_f, t) \Big|_{x_f=0} , \quad (3.10)$$

for each y_f . Let us mention that path integral representations of Green's functions for several types of differential equations satisfying local boundary conditions have already been studied extensively in more than one dimension (see Ref. [7] and references therein). The equation (3.9) was obtained in Ref. [7] for the diffusion equation. In the next section, we generalize this result for non-local boundary conditions.

3. The Functional Integral For General Boundary Condition

In this subsection, we construct a family of functional integrals $\mathcal{A}_g^B$ parameterized with the function g , introduced in Section II, which satisfies the boundary condition

$$\left. \frac{d}{dx_f} \mathcal{A}_g^B(\vec{z}_i, \vec{z}_f, \tau) \right|_{x_f=0} = \int dy'_f g(y'_f, y_f) \mathcal{A}_g^B(\vec{z}_i, (0, y'_f), \tau) \quad (3.11)$$

The analytic continuation of $\mathcal{A}_g^B$ with respect to τ results in the Green's functions (2.9) of the Schrödinger equation satisfying the non-local boundary condition (2.4) or equivalently (2.10).

In order to obtain a functional integral satisfying the non-local boundary conditions (3.11), we wish to define a measure giving non-zero weight to paths that can jump to other points on the edge of the half-plane when they hit the edge of the half-plane. Moreover, this measure should depend on the four functions ρ, α, f, h , introduced in Section II to parameterize the function g , since the boundary condition (3.11) depends on them. We will see below that when a path hits the edge of the half-plane at y along the y -axis, $\rho(y)$ gives the rate at which the path jumps from that point and $f(y, y')$ gives the density of probability of jumping to y' given it has jumped from y . The functions $\alpha(y)$ and $h(y, y')$ control the size of weights and phases given to the path when it spends time near y and when it jumps from y to y' respectively.

We will see that the probability of jumping from a point on the edge of the half-plane depends both on the time the path spends near that point and on the rate of jumping at that point. It is then natural to introduce the notion of *effective local time* which is roughly the amount of time the path spends near the edge of the half-plane weighted, point-by-point, by the rate at which the path jumps at that point. This notion is defined formally by (3.16) below.

Let us describe the process generating the paths on the half-plane. Imagine that the path starts at $\vec{z}_i \equiv (x_i, y_i)$, with $x_i > 0$ and undergoes ordinary reflected Brownian motion on the half-plane until its effective local time (defined below) reaches some number s_1 . (At the instant the effective local time reaches s_1 the path must be on the edge of the half-plane, say $(0, y_1^-)$.) It then jumps to $(0, y_1^+)$ on the edge of the half-plane. From this point it resumes its reflected Brownian motion on the half-plane until the effective local time reaches $s_1 + s_2$ at which time it reaches $(0, y_2^-)$ and jumps to $(0, y_2^+)$, and so forth.

The values $s_1, s_2, \dots$ are to be chosen as independent random variables which are exponentially distributed i.e.

$$\text{Prob}(s_k > u) = e^{-u} \quad (3.12)$$

As we will see later, the rate at which the paths jump is implicit in the definition of the effective local time, (3.16), and this is why we have set the coefficient of $-u$ to be 1 without loss of generality. The values $y_1^+, y_2^+, \dots$ are to be chosen as random variables and their distribution depends respectively on the locations $(0, y_1^-), (0, y_2^-), \dots$, on the edge of the half-plane, from which the jump takes place, that is

$$\text{Prob}(w < y_k^+ < w + dw) = f(y_k^-, w) dw \quad (3.13)$$

We see that $f(y_k^-, w)$ is the probability density of landing at $(0, w)$ given that the path has jumped from $(0, y_k^-)$.

More formally, let $\vec{q}^* = (q_x^*, q_y)$ be an ordinary reflected Brownian motion on the half-plane as described in Subsection III.1 and let ℓ be its local time defined via (3.3). Given the values $s_1, y_1^+, s_2, y_2^+, \dots$ as described above, we define the path $\vec{q}$ by

$$\vec{q}(\tau') \equiv (q_1(\tau'), q_2(\tau')) \equiv (q_x^*(\tau'), q_y(\tau') + \Delta y_1 + \Delta y_2 + \dots + \Delta y_{\#}) \quad (3.14)$$

where $\Delta y_k \equiv y_k^+ - y_k^-$, $\#$ is the number of jump,

$$s_1 + s_2 + \dots + s_{\#} \leq s(\tau') < s_1 + \dots + s_{\#+1} \quad (3.15)$$

and

$$s(\tau') \equiv \int_0^{\tau'} \rho(q_2(\tilde{\tau})) \dot{\ell}(\tilde{\tau}) d\tilde{\tau} \quad (3.16)$$

Here, $\rho(y)$ is the rate per unit local time at which the path jumps from the point $(0, y)$ on the edge of the half-plane. We call $s(\tau')$ the effective local time. It measures the “time” the path spends near the edge of the half-plane weighted, point-by-point, by the rate $\rho(y)$ at which the path jump at that point. Definitions (3.14) and (3.16) require further explanation since they are defined in terms of each other. Let (3.14) be given for the first $\#$ -jumps of the path, then (3.16) is well defined until $s(\tau')$ reaches $s_1 + s_2 + \dots + s_{\#+1}$. At this time the path is at $(0, y_{\#+1}^-)$ and jumps to $(0, y_{\#+1}^+)$. This determines $\Delta y_{\#+1}$ in definition (3.14), and (3.16) is updated to take this jump into account. $s(\tau')$ is now well defined until it reaches $s_1 + s_2 + \dots + s_{\#+2} \dots$ and so forth. The above is true for $\# = 0, 1, 2, \dots$ so definitions (3.14) and (3.16) should now be clear. Note that the values $y_1^-, y_2^-, \dots$ are completely determined by the half-plane reflected path $\vec{q}^*$ and the values $s_1, y_1^+, s_2, y_2^+, \dots$.

Let $\tau_1, \tau_2, \dots, \tau_{\#}$ be the times at which the effective local time reaches $s_1, s_1 + s_2, \dots, s_1 + s_2 + \dots + s_{\#}$. We note that $q_2(\tau')$ is discontinuous at $\tau_1, \tau_2, \dots, \tau_{\#}$. The positions of the path along the y -direction at time τ_k , before and after the k -th jump can be expressed in terms of $\vec{q} = (q_1, q_2)$ as

$$y_k^{\pm} \equiv \lim_{\epsilon \rightarrow 0} q_2(\tau_k \pm \epsilon) \quad (3.17)$$

The measure on paths $\vec{q}$, denoted as $[d\vec{q}]_{\rho, f} e^{-\frac{1}{2} \int_0^{\tau} \dot{q}^2 d\tau'}$, is induced from the measure on half-plane reflected paths and the distribution (3.12) and (3.13). We use this measure to define the functional integral

$$\mathcal{A}_g^B(\vec{z}_i, \vec{z}_f, \tau) = \int [d\vec{q}]_{\rho, f} e^{-\frac{1}{2} \int_0^{\tau} \dot{q}^2 d\tau'} e^{-\int_0^{\tau} \alpha(q_2(\tau')) \dot{\ell}(\tau') d\tau'} e^{i \sum_{k=1}^{\#} h(y_k^-, y_k^+)} \quad (3.18)$$

The functional integral is over all paths $\vec{q}$ from $\vec{z}_i \equiv (x_i, y_i)$ to $\vec{z}_f \equiv (x_f, y_f)$ in time τ . In (3.18), ρ, α, h, f are the four functions introduced in Section II in order to parameterize the function g specifying the boundary condition (2.4). $\#$ is the number of jumps of a given path $\vec{q}$. The last factor in (3.18), involving h , is a product of phases where each factor is the phase associated with one jump of the path. The arguments of h represent the location

from and to which the path jump respectively along the y -axis. The factor involving α was discussed in the previous subsection.

The memoryless property of the distribution (3.12) and the fact that the distribution of y_k^+ , (3.13), does not depend on past events but only on the present position of the path before a jump guarantees that $\mathcal{A}_g^B$ satisfies the convolution equation

$$\mathcal{A}_g^B(\vec{z}_i, \vec{z}_f, \tau_1 + \tau_2) = \int_{\text{half-plane}} d^2\vec{z} \mathcal{A}_g^B(\vec{z}_i, \vec{z}, \tau_1) \mathcal{A}_g^B(\vec{z}, \vec{z}_f, \tau_2) \quad (3.19)$$

(To establish this, we used the fact that (3.7) and the last factor in (3.18) convolve path-by-path). $\mathcal{A}_g^B$ also satisfies

$$\mathcal{A}_g^B(\vec{z}_i, \vec{z}_f, \tau) = \mathcal{A}_g^B(\vec{z}_f, \vec{z}_i, \tau)^* \quad (3.20)$$

because for each path $\vec{q}$ from $\vec{z}_i$ to $\vec{z}_f$ there exists a path $\vec{q}'$ from $\vec{z}_f$ to $\vec{z}_i$ with weight complex conjugate to the weight of the path $\vec{q}$. For $\vec{z}_i$ and $\vec{z}_f$ not on the edge of the half-plane, and for τ small (so that paths which hit the boundary can be neglected), $\mathcal{A}_g^B$ is well-approximated by $\mathcal{A}_{\text{wp}}^B$. Combined with (3.19) this ensures that $\mathcal{A}_g^B$ satisfies the same differential equation as $\mathcal{A}_{\text{wp}}^B$ for all τ and for all $\vec{z}_i, \vec{z}_f$ not on the edge of the half-plane:

$$-\frac{1}{2} \left(\frac{\partial^2}{\partial x_f^2} + \frac{\partial^2}{\partial y_f^2} \right) \mathcal{A}_g^B(\vec{z}_i, \vec{z}_f, \tau) = -\frac{d}{d\tau} \mathcal{A}_g^B(\vec{z}_i, \vec{z}_f, \tau) \quad (3.21)$$

with

$$\mathcal{A}_g^B(\vec{z}_i, \vec{z}_f, 0) = \delta(\vec{z}_f - \vec{z}_i) \quad . \quad (3.22)$$

The fact that $\mathcal{A}_g^B$ satisfies the last four equations is sufficient to guarantee it satisfies some boundary condition of the form (2.4). In the next subsection, we will show that indeed $\mathcal{A}_g^B$ satisfies the boundary condition (3.11). The analytic continuation $G_g(\vec{z}_i, \vec{z}_f, t) \equiv \mathcal{A}_g^B(\vec{z}_i, \vec{z}_f, it)$ satisfies (3.11) also. So G_g is the Green's function, (2.9), of the Schrödinger equation for the self-adjoint Hamiltonian corresponding to the boundary condition (2.4).

We can obtain the functional integral of more complicated quantum systems as follows. For a quantum system where the particle is limited to move inside a bounded region, one can show that the self-adjoint-extension of the Hamiltonian are indexed by a function g defined on the boundary of the region. For each self-adjoint-extension, we can construct the functional integral in terms of functions ρ, f, α, h , defined on the boundary of the region, parameterizing the function g . The functional integral we get is very similar to (3.18). The reflected Brownian measure needed in the definition of the functional integral is mathematically well-defined for region with smooth boundary and can be induced from the measure on whole line path.

So far in this paper, we have only considered the motion of a free particle. We can obtain the functional integral for a particle moving under the influence of a bounded potential, $U(\vec{q})$, by multiplying the integrand of the functional integral (3.18) by

$$e^{-\int_0^\tau U(\vec{q}(\tau')) d\tau'} \quad . \quad (3.23)$$

Both the self-adjoint extensions and the proof that the functional integral satisfies the appropriate boundary condition are unaffected by the introduction of that factor.

4. Proof of Boundary Condition for Functional Integral

In the previous subsection, we showed that the functional integral

$$\mathcal{A}_g^B(\vec{z}_i, \vec{z}_f, \tau) = \int [d\vec{q}]_{\rho, f} e^{-\frac{1}{2} \int_0^\tau \dot{\vec{q}}^2 d\tau'} e^{-\int_0^\tau \alpha(q_2(\tau')) \dot{\ell}(\tau') d\tau'} e^{i \sum_{k=1}^{\#} h(y_k^-, y_k^+)} \quad (3.24)$$

satisfies some boundary condition of the form (2.4). In this subsection, we prove that the functional integral satisfies the specific boundary condition

$$\left. \frac{d}{dx_f} \mathcal{A}_g^B(\vec{z}_i, \vec{z}_f, \tau) \right|_{x_f=0} = \int dy'_f g(y'_f, y_f) \mathcal{A}_g^B(\vec{z}_i, (0, y'_f), \tau) \quad (3.25)$$

We note that this boundary condition, regarded as a function of $\vec{z}_f$, is the same for each $(\vec{z}_i, \tau)$. Also, we remark that we only need to determine $\mathcal{A}_g^B$ near the edge of the half-plane and at a particular point along the y-axis in order to distinguish which boundary condition it satisfies among the different possibilities offered by (2.4). In other words, it is sufficient to analyze $\mathcal{A}_g^B$ in the limit

$$\left. \begin{aligned} \tau &\text{ small} \\ \vec{z}_i &\equiv (x_i, y_i) = (0, y_i) \\ \vec{z}_f &\equiv (x_f, y_f) \approx (0, y_f) \end{aligned} \right\} \quad (3.26)$$

where x_f is much smaller than $\sqrt{2\tau}$ and $y_f = y_i$. As the following shows, in this limit the path integral (3.24) is well-approximated by a sum over dominant paths. By Taylor expanding the last two factors in (3.24) around the dominant paths we see that to leading orders the Brownian motion in the y-direction decouples from the jumping process and the Brownian motion in the x-direction. Once the Brownian motion in the y-direction is integrated out, the path integral we are left with is recognized to be a good approximation for the path integral of a 1-dimensional quantum system made out of a continuous set of half-lines all joined together at the origin (see the Appendix). This identification enables us to use the techniques of Ref. [2] to show which boundary condition is satisfied by the path integral.

In the small τ -limit, the main contribution to the functional integral (3.24) comes from paths $\vec{q}(\tau')$ that remain within a circle of radius $\sqrt{2\tau}$ centered at (x_i, y_i) before the first jump and (x_i, y_k^+) between the k -th and $(k+1)$ -th jump. This is because between jumps the paths are generated by Brownian motion which have an exponentially small probability of reaching a point at a distance greater than $\sqrt{2\tau}$ from its starting point. Moreover, in the small τ -limit, paths for which $\ell(\tau)$ is greater than $\sqrt{2\tau}$ are exponentially suppressed^{2,6}. This implies, as shown in Ref. [2], that paths with $\#$ -jumps are suppressed by a factor $(2\tau)^{\# / 2}$. One can show that paths jumping less than three times contain all of the information necessary to determine which boundary condition is satisfied by $\mathcal{A}_g^B$. To establish this, it is sufficient to observe that, as a whole, these paths are sensible to all the values of the functions ρ, f, α, h : that is, any point on the boundary can be visited by at least one of these paths. If we had neglected paths that jump twice, this statement would not have been true. We can neglect paths that jump more than twice.

To sum up, we have seen that the dominant paths in the functional integral (3.24) jump less than three times, have a local time less than $\sqrt{2\tau}$ and remain within a circle of radius $\sqrt{2\tau}$ from the points (x_i, y_k^+) . (Note that $y_0^+ \equiv y_i$.) Also, we have seen that we can drop any contributions to the functional integral suppressed by at least a factor $(2\tau)^{3/2}$ since we have neglected paths with three jump and more.

In the functional integral (3.24) the functions ρ, f, α, h depend on the paths. Since a given path $\vec{q}$, with $\#$ -jump, deviates by at most $\sqrt{2\tau}$ from the points (x_i, y_k^+) , we can perform a Taylor expansion of ρ, f, α, h around these points. With a little bit of work, we can show that we need only to keep the leading terms in the expansion of these functions. For the contribution to the functional integral coming from the next to leading terms different from zero in this expansion are suppressed by a factor $(2\tau)^{3/2}$. (for the case where ρ, f, α, h are very slowly varying it is clear that it is sufficient to keep only the leading term on their expansion.)

By expanding ρ, f, α, h to leading order, we can rewrite the functional integral (3.24) as

$$\mathcal{A}_g^B(\vec{z}_i, \vec{z}_f, \tau) \approx \int_{\mathcal{D}} [d\vec{q}]_{\rho, f} e^{-\frac{1}{2} \int_0^\tau \dot{\vec{q}}^2 d\tau'} e^{-\sum_{k=0}^{\#} \alpha(y_k^+) \ell_k} e^{i \sum_{k=0}^{\#-1} h(y_k^+, y_{k+1}^+)} \quad (3.27)$$

where $\mathcal{D}$ stands for the path integral over all the dominant paths from $\vec{z}_i$ to $\vec{z}_f$; that is the paths that remain within a circle of radius $\sqrt{2\tau}$ from the points (x_i, y_k^+) , with zero, one and two jumps ($\# = 0, 1, 2$) and with local time less than $\sqrt{2\tau}$. (we have set $y_0^+ \equiv y_i$ and $y_k^- \approx y_{k-1}^+$, for $k = 1, 2, \dots$) Also,

$$\ell_k \equiv \lim_{\epsilon \rightarrow 0} \frac{1}{2\epsilon} \int_{\tau_k}^{\tau_{k+1}} I(q_1(\tau') < \epsilon) d\tau' \quad (3.28)$$

is the local time the path spends near the origin between the k -th and $(k+1)$ -th jump for $k = 1, 2, \dots, \# - 1$. ℓ_0 and $\ell_{\#}$ is the local time the path spends near the origin before the first and after the last jump respectively. Finally, the effective local time in the functional integral is

$$s(\tau) \approx \sum_{k=0}^{\#} \rho(y_k^+) \ell_k \quad (3.29)$$

to leading order in the expansion of ρ .

Now, let us make two remarks. First recall that a path $\vec{q}$ can be expressed in terms of a half-plane reflected path $\vec{q}^* = (q_x^*, q_y)$ from $\vec{z}_i$ to an intermediate point $(x_i, \tilde{y})$ and in terms of the values $s_1, y_1^+, s_2, y_2^+, \dots$ restricted to

$$\sum_{k=1}^{\#} \Delta y_k \equiv \sum_{k=1}^{\#} (y_k^+ - y_k^-) = y_f - \tilde{y} \quad (3.30)$$

This restriction ensures that the path $\vec{q}$ ends at $\vec{z}_f$ as required. The path integral can be broken into a few integrations. We begin by integrating over the values $s_1, y_1^+, s_2, y_2^+, \dots$ restricted to (3.30) with $\vec{q}^*$ and $\tilde{y}$ fixed. Next we integrate over paths q_x^*, q_y with $\tilde{y}$ fixed and finally integrate over all $\tilde{y}$. As a second remark we point out that, in the functional

integral (3.27), by relaxing the domain of integration to be over all paths instead of only those describe by $\mathcal{D}$ we only introduce errors suppressed by at least a factor $(2\tau)^{3/2}$.

We can now rewrite (3.27) as

$$\begin{aligned}\mathcal{A}_g^B(\vec{z}_i, \vec{z}_f, \tau) &\approx \int_{-\infty}^{+\infty} d\tilde{y} \mathcal{A}_{\text{free}}^B(y_i, \tilde{y}, \tau) \mathcal{A}_{y_i, y_i+(y_f-\tilde{y})}(x_i, x_f, \tau) \\ &\approx \mathcal{A}_{y_i, y_f}(x_i, x_f, \tau)\end{aligned}\tag{3.31}$$

where

$$\mathcal{A}_{\text{free}}^B(y_i, \tilde{y}, \tau) = \frac{1}{\sqrt{2\pi\tau}} e^{-(\tilde{y}-y_i)^2/2\tau}\tag{3.32}$$

is the result of the integration over all the paths q_y (note that to leading order nothing but $e^{-\frac{1}{2}\int_0^\tau \dot{q}^2 d\tau'}$ depends on $\vec{q}_y$ in (3.27).), and $\mathcal{A}_{y_i, y_f}$ is the result of the integration over the values $s_1, y_1^+, s_2, y_2^+, \dots$ restricted to (3.30) and over all paths q_x^* . In the appendix we show that $\mathcal{A}_{y_i, y_f}$ can be thought of as the functional integral of a 1-dimensional system made out of a continuous set of half-lines all joined together at the origin. This system is a straightforward generalization of a quantum system with only two half-lines connected at the origin and this system was studied in a previous paper². Using the same techniques introduced in Ref. [2] we can show that $\mathcal{A}_{y_i, y_f}$ satisfies

$$\left. \frac{d}{dx_f} \mathcal{A}_{y_i, y_f}(x_i, x_f) \right|_{x_f=0} = \int_{-\infty}^{+\infty} dy'_f g(y'_f, y_f) \mathcal{A}_{y_i, y'_f}(x_i, 0)\tag{3.33}$$

and in turn that implies that $\mathcal{A}_g^B$ satisfies (3.25) as we wanted to show.

IV. CONCLUSIONS

In this paper, the equivalence between the operator and functional approaches to quantum mechanics established for one-dimensional quantum systems in Ref. [1,2,3] has been extended to quantum systems in higher dimension admitting general boundary conditions. For each self-adjoint extension the Green's function is obtained as the analytic continuation of the functional integral whose measure depends explicitly upon the parameters defining the extension.

ACKNOWLEDGMENTS

I am grateful to Edward Farhi and to Sam Gutmann for their supports, insights and valuable remarks throughout the course of this work. I wish to thank Daniel Stroock, Barrett Rogers, Martin Leblanc and the referee for helpful discussions and comments.

REFERENCES

1. T.E. Clark, R. Menikoff and D. H. Sharpe, *Phys. Rev.* **D22** (1980) 3012.
E. Farhi and S. Gutmann, *Int. Jour. Mod. Phys.* **A5** (1990) 3029.
B. Gaveau and L. S. Shulman, *J. Phys. A.: Math. Gen.* **19** (1986) 1833.
2. M. Carreau, E. Farhi, S. Gutmann, *Phys. Rev.* **D42** (1990) 1194.
3. M. Carreau, E. Farhi, S. Gutmann and P. Mende, *Ann. of Phys.* **1**,(1990) 186.
4. K. Itô and H. McKean, Jr., *Diffusion Processes and Their Sample Paths* (Springer-Verlag, New York, 1964), p. 63.
5. Daniel Revuz, Marc Yor, *Continuous Martingales and Brownian Motion* (Springer-Verlag, New York, 1991) Chap. VI.
6. K. Sato and H. Tanaka, *Proc. Japan Acad.* **38** (1962) 699
7. K. Sato and T. Ueno, *J. Math. Kyoto Univ.* **4** (1965) 529
N. Ikeda, *Mem. Coll. Sci. Univ. Kyoto, Ser. A*, **33** (1961) 367

APPENDIX

Consider the quantum system made out of N -half-lines parameterized by $k = 1, 2, \dots, N$ all joined together at the origin as indicated in Fig.1. We locate the position of a particle on this system by specifying the branch, k , where the particle moves, and its distance, x , from the origin. The wave function $\chi_k(x)$ for each branch k of this system satisfies the Schrödinger equation. In order to make the Hamiltonian self-adjoint the wave function must satisfy the boundary condition

$$\left. \frac{d}{dx} \chi_j(x) \right|_{x=0} = \sum_{k=1}^N M_{k,j} \chi_k(0) \quad (A.1)$$

where M is a given hermitian matrix, characterizing a particular self-adjoint extension of this system. We can parameterize M in terms of the real parameters $\rho_j, \alpha_j, f_j, h_j$ as follows

$$M_{k,j} = (\rho_j + \alpha_j) \delta_{k,j} - \rho_k f_{k,j} e^{i h_{k,j}} \quad (A.2)$$

where $\rho_k \geq 0$, $f_{k,j} \geq 0$, $f_{k,k} = 0$, $h_{k,j} = -h_{j,k}$, $\rho_k f_{k,j} = \rho_j f_{j,k}$ and

$$\sum_{j=1}^N f_{k,j} = 1 \quad (A.3)$$

Below we will use these functions to construct the functional integral.

Let us describe the process generating the paths on this system. The path starts on branch i at x_i and undergoes reflected Brownian motion on this branch until its effective local time (defined below) reaches s_1 . At this time the path is at the origin and it jumps to branch i' at its origin and the path resumes its reflected Brownian motion until its effective local time reaches s_2 and so forth. The values $s_1, s_2, \dots$ are chosen to be independent random variables which are exponentially distributed, i.e.

$$\text{Prob}(s_k > u) = e^{-u} \quad (A.4)$$

Let $k_0, k_1, \dots, k_{\#}$ be the branch in which the path is moving before the first jump, after the first, second and last jump respectively. We choose the values $k_1, k_2, \dots$ to be random variables with the following distribution (note that $k_0 \equiv i$ is the branch in which the path starts its motion.) Given that the path has jumped from branch k_j , the distribution of the value k_{j+1} is

$$\text{Prob}(k_{j+1} = k') = f_{k_j, k'} \quad (A.5)$$

We see that $f_{k_j, k'}$ is the probability of jumping to branch k' given that the path had jumped from branch k_j . Let $\ell_0, \ell_1, \dots, \ell_{\#}$ be the local time the path spends near the origin in the branch k_0 before the first jump, and on branch $k_1, k_2, \dots, k_{\#}$ after the first, second, ..., last jump respectively, we define the effective local time as

$$s(\tau') = \sum_{j=0}^{\#} \rho_{k_j} \ell_j(\tau') \quad (A.6)$$

where $\#$ is the number of jumps of a given path and ρ_{k_j} is the rate per unit local time at which the paths jump from the origin of branch k_j .

More formally let q_x^* be a 1-dimensional reflected Brownian motion on the positive half-line. With $s_1, k_1, s_2, k_2, \dots$ as described above, we define the path $\bar{q}$ by

$$\bar{q}(\tau') = \left\{ q_x^*(\tau') \ ; \ \text{in branch } k_j \right\} \quad (A.7)$$

where

$$s_1 + s_2 + \dots + s_j \leq s(\tau') < s_1 + \dots + s_{j+1} \ . \quad (A.8)$$

The distribution (A.4) and (A.5) and the reflected Brownian measure for each path q_x^* induces a measure for each path $\bar{q}$. The functional integral for this system is

$$\mathcal{A}_{i,j}^B(x_i, x_f, \tau) = \int [d\bar{q}]_{\rho,f} e^{-\frac{1}{2} \int_0^\tau \dot{\bar{q}}^2 d\tau'} e^{-\sum_{j=0}^{\#} \alpha_{k_j} \ell_j} e^{i \sum_{j=0}^{\#-1} h(k_j, k_{j+1})} \quad (A.9)$$

where the path integral is over all paths $\bar{q}$ from x_i on branch i to x_f on branch j in time τ . Note that $k_0 \equiv i$, $k_{\#} \equiv j$ and that $e^{ih_{k,k'}}$ is a phase given to the path whenever it jumps from branch k to k' .

Equivalently the path integral can be performed by first integrating over all values $s_1, k_1, s_2, k_2, \dots$ restricted to

$$\sum_{j=1}^{\#} (k_j - k_{j-1}) \bmod N = j - i \quad (A.10)$$

with q_x^* fixed and then integrating over all paths q_x^* .

The functional integral (A.9) satisfies

$$\left. \frac{d}{dx_f} \mathcal{A}_{i,j}^B(x_i, x_f) \right|_{x_f=0} = \sum_{k=1}^N M_{k,j} \mathcal{A}_{i,k}^B(x_i, 0) \quad (A.11)$$

where M is given by (A.2). The proof of (A.11) is a straight forward generalization of the proof of the corresponding result for $N = 2$ given in Ref. [1] and will not be repeated here: for $N = 2$, (A.11) reduces to the Eq.(3.22) of Ref. [1] since hermiticity of the matrix M implies that $\rho_1 = \rho_2$, $f_{1,2} = f_{2,1} = 1$ and $h_{1,2} \equiv \theta$.

Now consider the quantum system made out of an infinite number of half-line parameterized by $-\infty < y < \infty$ all joined together at the origin. The path on this system starts at x_i on branch y_0 jumps to branch $y_1, y_2, \dots, y_{\#}$ whenever its effective local time reaches $s_1, s_2, \dots, s_{\#}$ respectively. We denote as $\ell_0, \ell_1, \dots, \ell_{\#}$ the local time the path spends near the origin before the first jump, after the second, third, ..., last jump respectively. The effective local time is defined as

$$s(\tau') = \sum_{k=0}^{\#} \rho(y_k) \ell_k(\tau') \quad (A.12)$$

where $\#$ is the number of jumps of a given path. The values $s_1, s_2, \dots$ are distributed as in (A.4). The values $y_1, y_2, \dots$ are distributed as follows. Given that the path has jumped from the origin of branch y_k , the distribution of the value y_{k+1} is

$$\text{Prob}(\nu < y_{k+1} < \nu + d\nu) = f(y_k, \nu) d\nu \quad . \quad (\text{A.13})$$

The paths $\bar{q}$ are defined as in (A.7) with k_j replace by y_j . The functional integral for this system is

$$\mathcal{A}_{y_i, y_f}^{\text{B}}(x_i, x_f, \tau) = \int [d\bar{q}]_{\rho, f} e^{-\frac{1}{2} \int_0^\tau \dot{\bar{q}}^2 d\tau'} e^{-\sum_{k=0}^{\#} \alpha(y_k) \ell_k} e^{i \sum_{k=0}^{\#-1} h(y_k, y_{k+1})} \quad (\text{A.14})$$

where the path integral is over all paths $\bar{q}$ from x_i on branch y_i to x_f on branch y_f in time τ . Note that $y_0 \equiv y_i$, $y_{\#} \equiv y_f$ and the function f, ρ, h and α are the same functions introduced in section II. Equivalently the path integral can be perform by first integrating over all values $s_1, y_1, s_2, y_2, \dots$ restricted to

$$\sum_{k=1}^{\#} (y_k - y_{k-1}) = y_f - y_i \quad (\text{A.15})$$

with q_x^* fixed and then integrating over all paths q_x^* . We can see now that the path integral $\mathcal{A}_{y_i, y_f}$ in (3.31) is given by the path integral (A.14). We can show that the functional integral (A.14) satisfies

$$\left. \frac{d}{dx_f} \mathcal{A}_{y_i, y_f}^{\text{B}}(x_i, x_f) \right|_{x_f=0} = \int_{-\infty}^{+\infty} dy'_f g(y'_f, y_f) \mathcal{A}_{y_i, y'_f}^{\text{B}}(x_i, 0) \quad (\text{A.16})$$

which is the generalization of (A.11).

Figure Captions

Fig. 1: Quantum system made out of N -half-lines joined together at the origin. As an example of a particle trajectory, a particle can start at x_i on branch 1 and can end up at x_f on branch 3.